

# Cover Crops in Vegetable Production Systems



CEREAL RYE

**C**urrent vegetable production systems require an intensive amount of work and inputs, and if not properly managed could have detrimental effects on soil and the environment. Practices such as intensive tillage, increased herbicide use, and reduced organic matter inputs add additional stress to the sustainability of vegetable production systems. Growers need the tools and best practices to make production systems sustainable without compromising farm productivity and profitability. Cover crops serve as a valuable production tool in preserving environmental sustainability of vegetable cropping systems and render numerous benefits to soil, vegetable crops, and the grower.

## **What is a cover crop?**

A cover crop is a crop that is not intended for harvest and is managed to maintain and improve soil fertility, water quality, and help manage weeds, pests, and diseases. Cover crops often are planted after harvesting a vegetable crop and then terminated before the planting of the next vegetable crop. There also are production systems where cover crops are used as living mulch, growing at the same time as the vegetable crop.

## **Benefits of cover crops**

Cover crops provide a wide range of ecological and environmental benefits. Depending on cover crop type and grower needs, each cover crop can be utilized to provide a specific ecological benefit. Table 1 provides a list of cover crops used in vegetable cropping systems. Some of the primary benefits which cover crop provide include:

### ***Soil and water conservation***

With the use of intensive tillage in vegetable production systems, there is a constant threat of soil erosion due to rain and wind. Cover crops prevent soil erosion by providing ground cover and plant roots to hold the soil.

Both the living foliage and the residue from dead cover crop plants protect the soil from rain drop impact and slow water and air flow across the soil surface, which reduces dislodging and movement of soil particles. The cover crop root system helps to hold soil in place by enmeshing and anchoring soil aggregates. Successive years of cover crop plantings can indirectly contribute to water conservation by

**SUSTAINABLE  
VEGETABLE  
PRODUCTION**

[www.extension.iastate.edu/vegetablelab](http://www.extension.iastate.edu/vegetablelab)

**TABLE 1. Cover crop characteristics for vegetable cropping systems**

Cover crop	Seeding rate* (lb/A)	Planting times	Expected biomass (lb/A)
<b>Brassicacae</b>			
Oilseed radish	10-12	Spring, late summer, early fall	4,000-6,000
Rapeseed	10-15	Fall, spring	2,000-4,000
Yellow mustard	10-12	Spring, late summer	4,000-8,000
<b>Legumes</b>			
Cowpeas	75-100	Early summer	2,500-5,000
Crimson clover	25-30	Early/late summer	3,500-5,000
Field peas	90-100	Fall, early spring	4,000-5,000
Hairy vetch	25-40	Early fall	2,500-5,000
Red clover	10-12	Early spring, late summer	2,000-5,000
Sweet clover	10-20	Spring	3,000-5,000
White clover	10-12	Spring, early fall	2,000-5,000
<b>Non-legumes</b>			
Annual ryegrass	20-30	Late summer, fall	3,000-5,000
Barley	75-125	Fall, spring	4,000-6,000
Buckwheat	50-80	Spring, late summer	2,000-5,000
Cereal rye	100-120	Late summer, fall	4,000-8,000
Oats	100-120	Early spring, late summer	4,000-8,000
Sorghum sudangrass	40-50	Late spring, early summer	8,000-12,000

\* Seeds broadcasted



increasing soil organic matter, which improves soil water holding capacity and infiltration. Successive years of cover crop plantings can indirectly contribute to water conservation by increasing soil organic matter, which improves soil water holding capacity and infiltration.

**Organic matter input**

A primary advantage of growing cover crops is the addition of organic matter to the soil. Organic matter improves the physical condition of the soil by improving the structure, aggregate stability, water

holding capacity, and porosity of the soil. Also, organic matter from cover crops improves nutrient cycling by increasing soil microbial population and activity. Examples of cover crops that can add substantial organic matter to soil include cereal rye, oats, sorghum-sudangrass, and triticale.

**Nitrogen fixation**

Leguminous cover crops such as clovers and vetches, have the added advantage of fixing atmospheric nitrogen for their growth and the following crops. This nitrogen fixation occurs through a symbiotic





**YELLOW CLOVER**



**YELLOW MUSTARD**

relationship between the leguminous plant and nitrogen-fixing bacteria that live in nodules (knobs) on the plant's roots. While the bacteria fix nitrogen for plant growth, the plant provides food and shelter to the bacteria. Upon death of the legume the nitrogen is released and around 40%–60% of the nitrogen in the legume cover crop is available to the next crop. The amount of nitrogen contributed by legumes varies by species (see Table 2). There are specific species of bacterium that form symbiotic relationship with individual legume cover crop species. It is advisable to inoculate legume seeds with the proper nitrogen-fixing bacterium strain for efficient nitrogen fixation. The cost for the inoculum packet is \$5–\$10 and can usually treat 50 pounds of seeds. Research has shown significant increase in cover crop biomass and nitrogen-fixing potential in inoculated legume cover crop systems.



**TABLE 2. Nitrogen fixation estimates by leguminous cover crops**

Cover crop	Nitrogen contributed (lb/A)
Crimson clover	50-125
Hairy vetch	100-120
Red clover	75-130
Sweet clover	50-125
White clover	80-130

**Nutrient scavenging**

Cover crops planted in the fall can scavenge and use unused soil nitrogen left at the end of the growing season, which may have otherwise leached during the fall or the spring. Certain cover crops tend to be very efficient at recycling or scavenging excess nutrients such as oilseed radish, cereal rye, yellow mustard, etc. These species are well adapted to cool, fall and spring conditions, and continue growing after nutrient absorption by the crop has slowed or stopped. When the cover crop dies, most of the nitrogen used by the plant during growth will be released and reused by future crops.

**Break soil compaction**

Cover crop roots can help alleviate the effects of soil compaction by penetrating a compacted layer and creating macropores or root channels that allow air, water and crop roots to penetrate deeper in the soil profile. Although all cover crop species contribute to loosening of soil, cover species differ in their capacity to penetrate compacted soils. In general, cover crops, such as oilseed radish, have large diameter taproots and are more effective at penetrating compacted soil layers than species that have small diameter roots. Once these taproots penetrate the restricting soil layer they are able to bring up nutrients from deep soil layers to upper layers of the soil.

**Enhance soil biology**

Soil is a living entity and is home to hundreds of thousands of different worms, insects, nematodes, and microorganisms. To keep soils healthy and improve soil quality, the value of cover crop root and shoot residues that help feed the soil throughout the entire year should be recognized.





The top 6 inches of soil can contain over 2,500 to 5,000 lbs./acre of living organisms. Cover crops improve the soil environment for both macro- and microorganisms, of which the majority are beneficial or not a problem for a vegetable crop. Cover crop residues increase soil organic matter, improve water holding capacity, provide a food source, and moderate soil temperature, all of which benefit soil macro- and microorganism communities. Several studies have shown higher soil microbial biomass and diverse soil microbial populations under cover-cropped systems. Cover crops also promote populations of soil macrofauna such as earthworms, millipedes, beetles, and spiders, which help create air pore spaces in the soil.

### ***Bio-fumigation***

Cover crops can be used to suppress problematic plant pathogenic nematodes, bacteria, and fungi in the soil. Certain cover crops in the Brassicaceae family (plants with cross-shaped petals) produce biologically active compounds, called glucosinolates, that have shown activity on soil-borne pests. Glucosinolates are present in plant roots, shoots, stems, and leaves and when incorporated into the soil they break down into compounds called isothiocyanates (ITCs) and other chemicals. The ITCs are known to suppress soil-borne diseases, nematodes, and weed seeds. Some cover crops that belong to the Brassicaceae family include oilseed radish, canola, Indian mustard, brown mustard, and yellow mustard. It is important to

note that these cover crops cannot be used as a sole control measure to mitigate soil pest problems; rather they should be used to enhance management strategies. Additionally, there is variability in the biofumigation capabilities, a technique of incorporating a plant's biomass into the soil, which will release toxic volatiles that suppress pests, among varieties of cover crops. For example, oilseed radish cultivars such as Adagio and Ultimo which have European origin, are reported to give better nematode suppression (especially cyst nematodes) than other cultivars. Oilseed radish cultivars commercially available and commonly grown in United States include Defender and Daikon.

### ***Weed suppression***

Cover crops can be used to manage weeds in vegetable production systems. Cover crops can reduce weed germination and establishment by competing and/or producing allelochemicals, which suppress weed seed germination. Cover crops such as cereal grains and grasses establish quickly in the fall, cover the soil, and grow throughout the winter, thereby suppressing fall and winter weeds. Small-seeded legumes that are seeded in the fall are sometimes not a good choice for weed suppression as they grow slowly during cold weather and can be outcompeted by weeds. Cover crops can influence weeds either in the form of living plants or as plant residue remaining after the cover crop is killed.





**HAIRY VETCH**



**INSPECTING ROOTS**

**Crop rotation**

Crop rotation is a planned system of growing different crops in succession on the same land. Benefits of crop rotation in terms of weed, pest, and disease management are well documented. Cover crops can be used in crop rotation plans to break pest cycles, add organic matter, and improve soil quality and health. Vegetables have many potential seasons of production, and given the choices available with long- and short-term cover crop life cycles, cover crops can easily fit into any crop rotation plan. Periods of 1–2 months between harvest of early planted spring crops and planting of fall crops can be filled using fast-growing, warm-season cover crops, such as buckwheat, cowpea, oats, and sorghum-sudangrass. Table 3 (page 6) provides a few examples and scenarios of how cover crops could be integrated with vegetable cropping systems.



**COVER CROP FIELD DAY**



**CLASSROOM WORKSHOP**



No endorsement is intended by Iowa State University Extension and Outreach of companies or their products mentioned nor is criticism implied of similar companies or their products not mentioned.





## Conclusion

Cover crops are gaining importance and are becoming an integral part of vegetable cropping systems. They improve the sustainability of vegetable production systems by reducing soil erosion, compaction and synthetic nitrogen inputs, suppressing weeds, increasing soil organic matter and water infiltration, enhancing soil biology, and providing habitat for beneficial insects and natural enemies of pests.

## Resources

Kaspar, T.C. and J.W. Singer. 2011. "The use of cover crops to manage soil," p. 321-337, In: J. L. Hatfield and T. J. Sauer (eds.). *Soil management: Building a stable base for agriculture*. American Society of Agronomy and *Soil Science Society of America Journal*, Madison, Wis.

*Managing Cover Crops Profitably*, 3rd ed. 2012. Sustainable Agriculture Research and Education (SARE), National Institute of Food and Agriculture, U.S. Department of Agriculture. Accessed May 1, 2015. <http://sare.org/publications/covercrops.htm>

Michigan Cover Crops website, Michigan State University Extension Agriculture and Agri-business Institute, East Lansing, Mich. Accessed May 1, 2015. <http://covercrops.msu.edu/>

Ngouajio, M. and D. Mutch. 2004. "Oilseed Radish: A New Cover Crop for Michigan," Bulletin-E2907, Michigan State University Extension Agriculture and Agri-business Institute, East Lansing, Mich.

Mohler, C. L. and S. E. Johnson, (eds.) 2009. "Crop Rotation on Organic Farms: A Planning Manual." NRAES 177. Sustainable Agriculture Research and Education (SARE), Plant and Life Sciences Publishing.



This product was developed with support from the Sustainable Agriculture Research and Education (SARE) program, which is funded by the U.S. Department of Agriculture—National Institute of Food and Agriculture (USDA-NIFA). Any opinions, findings, conclusions or recommendations expressed within do not necessarily reflect the view of the SARE program or the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.



Prepared by Ajay Nair, assistant professor of horticulture and extension vegetable production specialist Iowa State University, Ames, Iowa; Tom Kaspar, plant physiologist, USDA-ARS, National Laboratory for Agriculture and Environment, Ames, Iowa; and Gail Nonnecke, Morrill professor and global professor of horticulture, Iowa State University, Ames, Iowa.

Photo Credits:

Ajay Nair

Page 3, crimson clover illustration

(Source: Britton, N.L., and A. Brown. 1913.

An illustrated flora of the northern United States, Canada and the British Possessions. 3 vols.

Charles Scribner's Sons, New York. Vol. 2: 355.

Page 5, buckwheat illustration by Spline\_x (iStock.com)

**IOWA STATE UNIVERSITY**  
Extension and Outreach

[www.extension.iastate.edu](http://www.extension.iastate.edu)

### ... and justice for all

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotope, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue SW, Washington, DC 20250-9410, or call 800-795-3272 (voice) or 202-720-6382 (TDD). USDA is an equal opportunity provider and employer.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Cathann A. Kress, director, Cooperative Extension Service, Iowa State University of Science and Technology, Ames, Iowa.